

# AmpliBoron™

MIX WELL  
BEFORE USE

ACT 36, 1947 B3937

Trace element concentrate to correct and prevent deficiencies in agriculture cropping. Boron is a crucial trace element in maintaining normal physiological functions.

Nitrogen	(N)	40	g/kg
Boron	(B)	140	g/kg

**APPLICATION GUIDE:** Apply through Boomspray, aerial or overhead irrigation. DO NOT apply when conditions are extremely hot. Best applied in temperatures in the range of 5 – 25°C.

#### MIXING INSTRUCTIONS

1. Do not apply as a concentrate with other chemicals.
2. Half fill tank to the required volume with water.
3. Add the required amount of **AmpliBoron™**.
4. Add appropriate amount of non-ionic wetting agent as per manufacturer's instructions.
5. Fill the remainder of tank with water and maintain agitation.

#### STORAGE

DO NOT STORE product in direct sunlight.

DO NOT STORE product in temperature below 5°C or above 30°C.

DO NOT STORE DIRECTLY ON FLOOR, product to be raised from ground with pallettes.

#### COMPATIBILITY

Product is compatible with majority of agro chemicals, please check with the manufacturer. Please be aware that physical compatibility does not guarantee efficacy. Jar test before use.

#### CONDITIONS OF SALE

These goods are manufactured and tested to specific standards and sold in good faith, however, no guarantee is given due to the failure by the user to follow the labels instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration.

**KEEP OUT OF REACH  
OF CHILDREN.**

SPECIFIC GRAVITY  
1.38

#### NETT CONTENTS:

VOLUME	WEIGHT
500ml	0.69 kg
1L	1.38 kg
5L	6.9 kg
20L	27.6 kg
25 L	34.5 kg
200L	276 kg
1000L	1380 kg

BROADACRE	RATE L/Ha	WATER	COMMENTS
Cotton	1 - 2	1 : 100	Apply prior to flowering & again as boll fills
Wheat	1 - 2	1 : 100	Apply prior to flowering
Legumes	1 - 2	1 : 100	Apply 3-4 weeks after emergence
Canola	1 - 2	1 : 100	Apply prior to flowering
Maize/Sorghum	1 - 2	1 : 100	Apply prior to flowering

VITICULTURE	RATE L/Ha	WATER	COMMENTS
Wine & Table Grapes	1 - 1.5	1 : 500	3 Applications required 1. Prior to flowering 2. Flower buds separate 3. Fruit set

HORTICULTURE	RATE L/Ha	WATER	COMMENTS
Beans	1 - 2	1 : 200	Apply at 4-6 true leaf stage
Brassica	1 - 3	1 : 200	Apply prior to flowering & at stem extension
Cucurbits	1 - 3	1 : 200	Apply at 4-6 true leaf and prior to flowering
Carrots	1 - 2	1 : 200	Apply at seedling and vegetative stage
Lettuce	1 - 1.5	1 : 200	Apply at 4-6 leaf
Potatoes	1 - 2	1 : 200	Apply at seedling and prior to flowering
Strawberries	1 - 2	1 : 300	Apply prior to flowering
Tomatoes	1 - 2	1 : 200	Apply at seedling and prior to flowering

TREE CROPS	RATE L/Ha	WATER	COMMENTS
Stone Fruit	1 - 1.5	1 : 500	Apply at pink bud & again at after petal fall
Pome Fruit	1 - 1.5	1 : 500	Apply at spur leaves unfolded & after petal fall
Citrus	1 - 2	1 : 1000	Apply prior to flowering
Avocado	2 - 3	1 : 500	Apply Prior to flowering and after fruit set
Nut trees	1 - 2	1 : 500	Apply at spring flush to flowering and post harvest
Olives	1 - 2	1 : 300	Apply prior to flowering



Sustainable Soil Solutions



Contact us for more information: **Head Office:** (018) 293 3530 • **Central Region:** (056) 212 9565  
• **Cape Region:** (021) 555 2310 • **Natal Region:** (032) 486 1571 • **Eastern Cape:** (041) 486 2365  
e-mail: [info@profert.co.za](mailto:info@profert.co.za) [www.profert.co.za](http://www.profert.co.za)