

AmpliPhos™

MIX WELL
BEFORE USE

9.13.0. (22)

ACT 36, 1947 K8136

A concentrated neutral polyphosphate solution complexed with organic catalysts that can be used as a starter and foliar fertiliser. Unique formulation which support the energy system in plants and availability of phosphate in soils.

Nitrogen (N) 90 g/kg
Phosphate (P) 130 g/kg

APPLICATION GUIDE: Apply through Boomspray, aerial or overhead irrigation. DO NOT apply when conditions are extremely hot. Best applied in temperatures in the range of 5 – 25°C.

MIXING INSTRUCTIONS

1. Do not apply as a concentrate with other chemicals.
2. Half fill tank to the required volume with water.
3. Add the required amount of **AmpliPhos™**.
4. Add appropriate amount of non-ionic wetting agent as per manufacturer's instructions.
5. Fill the remainder of tank with water and maintain agitation.

STORAGE

DO NOT STORE product in direct sunlight.

DO NOT STORE product in temperature below 5°C or above 30°C.

DO NOT STORE DIRECTLY ON FLOOR, product to be raised from ground with pallets.

COMPATIBILITY

Product is compatible with majority of agro chemicals, please check with the manufacturer. Please be aware that physical compatibility does not guarantee efficacy. Jar test before use.

CONDITIONS OF SALE

These goods are manufactured and tested to specific standards and sold in good faith, however, no guarantee is given due to the failure by the user to follow the labels instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration.

SPECIFIC GRAVITY
1.38

NETT CONTENTS:
VOLUME WEIGHT
500ml 0.69 kg
1L 1.38 kg
5L 6.9 kg
20L 27.6 kg
25 L 34.5 kg
200L 276 kg
1000L 1380 kg

KEEP OUT OF REACH
OF CHILDREN.

BROADACRE	RATE L/Ha	WATER	COMMENTS
Cotton			
- Soil Injection	5 - 12	1 : 10 - 20	Apply at plant with water injection
- Foliar	4 - 10	1 : 50	Apply as foliar at 6 - 8 leaf stage
Wheat			
- Soil Injection	25 - 60	1 : 20 - 40	Apply at plant for P retentive soil
- Foliar	4 - 10	1 : 50	Apply as foliar at early tillering
Legumes	4 - 10	1 : 50	Apply 3-4 weeks after emergence
Canola	4 - 10	1 : 50	Apply 4 - 6 leaf to early cabbage
Maize/Sorghum	4 - 10	1 : 50	Apply 4 - 6 leaf up to flowering

HORTICULTURE	RATE L/Ha	WATER	COMMENTS
Beans	3 - 5	1 : 100	Apply at 4-6 true leaf stage
Brassica	3 - 5	1 : 100	Apply at 4-6 leaf stage
Cucurbits	3 - 5	1 : 100	Apply at seedling and prior to flowering
Carrots	3 - 5	1 : 100	Apply at seedling and vegetative stage
Lettuce	3 - 5	1 : 100	Apply at 4-6 leaf
Potatoes	3 - 5	1 : 100	Apply at seedling and prior to flowering
Strawberries	3 - 5	1 : 100	Apply at 4-6 leaf stage
Tomatoes	3 - 5	1 : 100	Apply at seedling and prior to flowering

TREE CROPS	RATE L/Ha	WATER	COMMENTS
Citrus	5 - 10	1 : 400	Apply during flush
Avocado	5 - 10	1 : 200	Apply during flush to flowering
Olives	5 - 10	1 : 200	Apply prior to flowering
Fertigation	10 - 20		Apply 2 - 3 applications prior to flowering



Sustainable Soil Solutions



Contact us for more information: Head Office: (018) 293 3530 • Central Region: (056) 212 9565
• Cape Region: (021) 555 2310 • Natal Region: (032) 486 1571 • Eastern Cape: (041) 486 2365
e-mail: info@profert.co.za www.profert.co.za